

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PIPERACILLIN and ZOSAMOX (Injection) safely and effectively. See full prescribing information for PIPERACILLIN AND ZOSAMOX FOR INJECTION.

PIPERACILLIN AND ZOSAMOX (Injection) is a combination product for intravenous use - Pharmacy Bulk Package Bottle Initial U.S. approval: 1993

PHARMACY BULK PACKAGE - NOT FOR PATIENT USE RECENT MAJOR CHANGES

Warnings and Precautions, Hemophagocytic Lymphohistiocytosis (5.3)

INDICATIONS AND USAGE

Piperacillin and tazobactam is a combination of piperacillin, a penicillin-class antibiotic and tazobactam, a beta-lactamase inhibitor, indicated for the treatment of:

- Intracranial pneumonia in adult and pediatric patients 2 months of age and older (1.1)
Nosocomial pneumonia in adult and pediatric patients 2 months of age and older (1.2)
Skin and skin structure infections in adults (1.3)
Serious respiratory infections in adults (1.4)
Community-acquired pneumonia in adults (1.5)
reduce the development of drug-resistant bacteria and maintain the effectiveness of piperacillin and tazobactam for injection against both aerobic gram-negative, piperacillin and tazobactam for injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. (1.6)

DOSEAGE AND ADMINISTRATION

Adult Patients with Indications Other Than Nosocomial Pneumonia. The recommended dosage is a dosage of 4.5 every six hours plus amoxicillin 3.375 g every six hours totaling 13.5 g (12 g piperacillin and 1.5 g tazobactam) every 6 hours (2.1)

Adult Patients with Nosocomial Pneumonia. Initial presumptive treatment of patients with nosocomial pneumonia should start with piperacillin and tazobactam for injection at a dosage of 4.5 every six hours plus amoxicillin, totaling 18 g (16 g piperacillin and 2 g tazobactam). (2.2)
Adult Patients with Renal Impairment. Patients with renal impairment (creatinine clearance < 40 mL/min) and dialysis patients should be reduced, based on the degree of renal impairment. (2.3)

Recommended Dosage of Piperacillin and Tazobactam for Pediatric Patients 2 months of Age and Older, Weighing up to 40 kg and With Normal Renal Function

Table with 3 columns: Age, Appendicitis and/or Peritonitis, Nosocomial Pneumonia. Rows for 2 months to 9 months and Older than 9 months.

Administer piperacillin and tazobactam for injection by intravenous infusion over 30 minutes. (2.4)

Contraindications. Do not administer piperacillin and tazobactam for injection to patients with known hypersensitivity to piperacillin and tazobactam for injection or to any of the components of the formulation. (2.5)

Warnings and Precautions. Hemophagocytic Lymphohistiocytosis (5.3)
Hypersensitivity Adverse Reactions (5.1)
Severe Cutaneous Adverse Reactions (5.6)
Neutropenia in Critically Ill Patients (5.5)
Clostridioides difficile-Associated Diarrhea (5.4)
Development of Drug-Resistant Bacteria (5.2)

Additional Experience with Piperacillin

Compatibility With Aminoglycosides

DOSEAGE FORMS AND STRENGTHS

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

WARNINGS AND PRECAUTIONS

INDICATIONS AND USAGE

DOSEAGE AND ADMINISTRATION

CONTRAINDICATIONS

DOSEAGE FORMS AND STRENGTHS

Piperacillin and tazobactam for injection: 40 x 5 lyophilized powder for reconstitution in pharmacy bulk package bottle. (3)

CONTRAINDICATIONS

Patients with a history of allergic reactions to any of the penicillins, cephalosporins, or beta-lactams inhibitors. (4)

WARNINGS AND PRECAUTIONS

Serious hypersensitivity reactions (anaphylactic/anaphylactoid) reactions have been reported in patients receiving piperacillin and tazobactam for injection. Discontinue piperacillin and tazobactam for injection if a reaction occurs. (5.1)

Piperacillin and tazobactam for injection may cause severe cutaneous adverse reactions, such as Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis. Discontinue piperacillin and tazobactam for injection for progressive rashes. (5.2)

Hemophagocytic lymphohistiocytosis (HLH) has been reported with the use of piperacillin and tazobactam for injection. If HLH is suspected, Hematologic effects (including bleeding, leukopenia and neutropenia) have occurred. Monitor hematologic tests during prolonged therapy. (5.3)
Neutropenia in critically ill patients was associated with the use of piperacillin and tazobactam for injection was found to be an independent risk factor for renal failure and was associated with delayed recovery of renal function as compared to other beta-lactam antibiotics used in a randomized, multicenter, controlled trial in critically ill patients. (5.4)
Neutropenia in critically ill patients was associated with the use of piperacillin and tazobactam for injection was found to be an independent risk factor for renal failure and was associated with delayed recovery of renal function as compared to other beta-lactam antibiotics used in a randomized, multicenter, controlled trial in critically ill patients. (5.5)

Neutropenia in critically ill patients was associated with the use of piperacillin and tazobactam for injection was found to be an independent risk factor for renal failure and was associated with delayed recovery of renal function as compared to other beta-lactam antibiotics used in a randomized, multicenter, controlled trial in critically ill patients. (5.6)

ADVERSE REACTIONS

The most common adverse reactions (incidence > 5%) are diarrhea, constipation, nausea, headache, and insomnia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Piperacillin and tazobactam for injection administration can significantly reduce the activity of aminoglycosides in hemodialysis patients. Monitor tuberculin concentrations in these patients. (7.1)
Probenecid prolongs the half-life of piperacillin and tazobactam and should not be co-administered with piperacillin and tazobactam for injection unless the benefit outweighs the risk. (7.2)
Co-administration of piperacillin and tazobactam for injection with vancomycin may increase the incidence of acute kidney injury. Monitor kidney function in patients receiving piperacillin and tazobactam for injection and vancomycin. (7.3)
Monitor coagulation parameters in patients receiving piperacillin and tazobactam for injection. (7.4)
Piperacillin and tazobactam for injection may prolong the neuromuscular blockade of vecuronium and other non-depolarizing neuromuscular blockers. (7.5)

USE IN SPECIFIC POPULATIONS

Dosage in patients with renal impairment (creatinine clearance < 40 mL/min) should be reduced based on the degree of renal impairment. (2.3, 8.6)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2024

7 DRUG INTERACTIONS

- 7.1 Aminoglycosides
7.2 Probenecid
7.3 Vancomycin
7.4 Anticoagulants
7.5 Vecuronium
7.6 Neuroleptics
7.7 Effects on Laboratory Tests

8 USE IN SPECIFIC POPULATIONS

- 8.1 Lactation
8.2 Pediatric Use
8.3 Geriatric Use
8.4 Renal Impairment
8.5 Hepatic Impairment
8.6 Patients with Cystic Fibrosis

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.2 Pharmacokinetics
12.3 Pharmacodynamics
12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

Sections or subsections omitted from the full prescribing information are not listed.

Table 1: Recommended Dosage of Piperacillin and Tazobactam for Injection in Patients with Normal Renal Function (As Total grams piperacillin and tazobactam\*)

Table with 3 columns: Creatinine clearance, mL/min; Greater than 40 mL/min; 30-40 mL/min; 15-30 mL/min; Less than 20 mL/min\*\*

\* CAPD 2.25 every 12 hours; 2.25 every 8 hours

\*\* 0.75 g of piperacillin and 0.08 g tazobactam should be administered every two hours for all indications other than nosocomial pneumonia and 2.25 g every eight hours for nosocomial pneumonia. Since amoxicillin is not included in the combination product, the total dosage of additional dose of 0.75 g piperacillin and tazobactam for injection (0.57 g piperacillin and 0.08 g tazobactam) should be administered following each dialysis period on hemodialysis days. No additional dosage of piperacillin and tazobactam for injection is necessary for CAPD patients.

For patients on hemodialysis, the maximum dose is 2.25 g every two hours for all indications other than nosocomial pneumonia and 2.25 g every eight hours for nosocomial pneumonia. Since amoxicillin is not included in the combination product, the total dosage of additional dose of 0.75 g piperacillin and tazobactam for injection (0.57 g piperacillin and 0.08 g tazobactam) should be administered following each dialysis period on hemodialysis days. No additional dosage of piperacillin and tazobactam for injection is necessary for CAPD patients.

2.4 Dosage in Pediatric Patients With Appendicitis/Peritonitis or Nosocomial Pneumonia
Piperacillin and tazobactam for injection for pediatric patients with appendicitis and/or peritonitis or nosocomial pneumonia aged 2 months of age and older should weigh up to 40 kg, and with normal renal function, should be administered as follows: (See Use in Specific Populations (8.4) and Clinical Pharmacology (12.3).)

Table 2: Recommended Dosage of Piperacillin and Tazobactam for Injection in Pediatric Patients with Normal Renal Function\*

Table with 3 columns: Age, Appendicitis and/or Peritonitis, Nosocomial Pneumonia. Rows for 2 months to 9 months and Older than 9 months of age.

\* Administer piperacillin and tazobactam for injection by intravenous infusion over 30 minutes.

Pediatric patients weighing over 40 kg and with normal renal function should receive the adult dose. (See Dosage and Administration (2.1, 2.2).)

Dosage of piperacillin and tazobactam for injection in pediatric patients aged 2 months to 9 months should be reduced based on the degree of renal impairment. (2.3)

2.5 Reconstitution and Dilution of Piperacillin and Tazobactam for Injection

Reconstitution of Piperacillin and Tazobactam for Injection for Adult Patients with Indications Other Than Nosocomial Pneumonia

Pharmacy bulk package bottles
Reconstituted pharmacy bulk package bottle solution will be transferred and further diluted for intravenous infusion.

The pharmacy bulk package bottle is for use in a hospital pharmacy admixture service only under a laminar flow hood. After reconstitution, the pharmacy bulk package bottle should be made with a sterile transfer set or other sterile dispensing device, and contents should be dispensed as needed into intravenous solution using aseptic technique. The contents of pharmacy bulk package bottle promptly. Discard unused portion after 24 hours if stored at room temperature (20°C to 25°C [68°F to 77°F]) or 48 hours if stored at refrigerated temperature (2°C to 8°C [36°F to 46°F]).

Reconstitute the pharmacy bulk package bottle with exactly 152 mL of sterile water for injection to produce a reconstituted solution containing 200 mg/mL of piperacillin and 25 mg/mL of tazobactam. Shake well

until dissolved. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to and during administration. Do not use if the solution is cloudy or contains any undissolved particles. Compatible Reconstitution Diluents for Pharmacy Bulk Package Bottles: 0.9% sodium chloride for injection

0.9% sodium chloride for injection
Dextrose 5%
Bactericidal water/parabens
Bactericidal water/benzyl alcohol
Bactericidal water/benzyl alcohol

Dilution of the Reconstituted Piperacillin and Tazobactam for Injection Solution for Adult Patients and Pediatric Patients Weighing Over 40 kg
Reconstituted piperacillin and tazobactam for injection pharmacy bulk package bottles should be further diluted (reconstituted) in 50 mL or 150 mL of 0.9% sodium chloride for injection intravenous solution listed below. Administer by infusion over a period of at least 30 minutes. During the infusion it is desirable to discontinue use of any other intravenous solution.
Compatible Intravenous Solutions for Pharmacy Bulk Package Bottles: 0.9% sodium chloride for injection
Sterile water for injection
Lactated Ringers solution
Dextrose 5% in saline

CAUTION: PIPERACILLIN SOLUTION IS NOT COMPATIBLE WITH PIPERACILLIN AND ZOSAMOX FOR INJECTION.
Piperacillin and tazobactam for injection should not be mixed with other drugs in a syringe or infusion bottle since compatibility has not been established.

Piperacillin and tazobactam for injection is not chemically stable in solutions that contain only sodium bicarbonate and solutions that contain only sodium chloride.
Piperacillin and tazobactam for injection should not be added to blood products or albumin hydrolysates. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Dilution of the Reconstituted Piperacillin and Tazobactam Solution for Pediatric Patients Weighing Up to 40 kg
The volume of reconstituted solution required to deliver the dose of piperacillin and tazobactam for injection to a child is dependent on the weight of the child. (See Dosage and Administration (2.4)). Reconstituted Piperacillin and Tazobactam solutions for pharmacy bulk package bottles should be further diluted in a compatible intravenous solution listed above.

- 1. Calculate patient dose as described in Table 2 above (See Dosage and Administration (2.4)).
2. Reconstitute bottle with a compatible reconstitution diluent, as listed in Table 1 above (See Dosage and Administration (2.4)).
3. Add the reconstituted solution to the Pharmacy Bulk Package Bottles, using the appropriate volume of diluent, as listed in table 4 below. Following the addition of the reconstitution diluent, the Pharmacy Bulk Package bottle until the powder is completely dissolved.

Table 4: Reconstitution of Pharmacy Bulk Package Bottle and Dilution Concentration

Table with 3 columns: Strength per Pharmacy Bulk Package Bottle, Volume of Diluent to be Added to the Bottle, Concentration of the Reconstituted Solution. Rows for 40.5 g/152 mL and 4.5 g/15.2 mL.

- 3. Calculate the required volume (mL) of reconstituted piperacillin and tazobactam for injection to deliver the patient dose.
4. Aseptically withdraw the required volume of reconstituted piperacillin and tazobactam solution from the pharmacy bulk package bottle. The volume should be further diluted in 50 mL or 150 mL of 0.9% sodium chloride for injection intravenous solution listed above.
5. Administer the diluted piperacillin and tazobactam for injection solution by infusion over a period of at least 30 minutes (a programmable syringe or infusion pump is recommended). During the infusion it is desirable to discontinue the primary infusion solution.

Stability of Piperacillin and Tazobactam for Injection Following Reconstitution

Piperacillin and tazobactam for injection reconstituted from pharmacy bulk package bottles are stable in glass and plastic containers (patients with renal impairment) for up to 24 hours at room temperature. The pharmacy bulk package bottle should NOT be frozen after reconstitution.

Pharmacy bulk package bottles should be used immediately after reconstitution. Discard any unused portion of the solution if stored at room temperature (20°C to 25°C [68°F to 77°F]), or after storage at 2°C to 8°C (36°F to 46°F) for up to 48 hours (refrigerated). (See 7.5.)

Stability studies in the I.V. bags have demonstrated chemical stability (potency, pH) of reconstituted solution and dilution of solution for up to 24 hours at room temperature and for up to 48 hours at refrigerated temperatures. Piperacillin and tazobactam for injection contains no preservatives. Appropriate consideration of aseptic technique should be used.

Piperacillin and tazobactam for injection reconstituted from pharmacy bulk package bottles can be used in ambulatory intravenous infusion pumps. Stability of piperacillin and tazobactam for injection in ambulatory intravenous infusion pumps has been demonstrated for a period of 24 hours at room temperature. Each 100 mL of reconstituted solution diluted to a volume of 37.5 mL or 25 mL. One-day supply of dosing solution were aseptically transferred into this medication reservoir (reservoir 100 mL or 250 mL) and stored for up to 24 hours in an ambulatory intravenous infusion pump per the manufacturer's instructions. Stability of piperacillin and tazobactam for injection in an infusion pump was not affected when administered using an ambulatory intravenous infusion pump.

2.6 Compatibility with Aminoglycosides

Due to the in vitro inactivation of aminoglycosides by piperacillin, piperacillin and tazobactam for injection and aminoglycosides should be administered separately. If separate administration of piperacillin and tazobactam for injection and aminoglycosides should be reconstituted, the order of administration is important when concomitant therapy with aminoglycosides is indicated. (See Drug Interactions (7.1).)

In circumstances where co-administration via Y-site is necessary, piperacillin and tazobactam for injection should be reconstituted for simultaneous co-administration via Y-site infusion with the following aminoglycosides under the following conditions:

Table with 3 columns: Aminoglycoside, Piperacillin and Tazobactam for Injection Concentration, Aminoglycoside Concentration Range (mg/mL), Acceptable Diluent. Rows for Amikacin, Gentamicin.

\* Diluent volumes apply only to bulk pharmacy package bottles
† Concentration ranges in Tables 4 and 5 are based on administration of the aminoglycoside in divided doses (10 - 15 mg/kg IV in daily doses for amikacin and 3 - 5 mg/kg IV in three daily doses for gentamicin) over 24 hours. The concentration of aminoglycoside in these states above via Y-site with piperacillin and tazobactam for injection has not been evaluated. Intermittent bolus dosing of aminoglycosides is not recommended.

Only the concentration and diluents for amikacin or gentamicin with the dosages of piperacillin and tazobactam for injection listed above are recommended for simultaneous co-administration via Y-site infusion. Simultaneous co-administration via Y-site infusion in any other diluent than those listed above is not recommended. The compatibility of the aminoglycosides by piperacillin and tazobactam for injection.

Piperacillin and tazobactam for injection is not compatible with other drugs for simultaneous co-administration via Y-site infusion.
Compatibility of piperacillin and tazobactam for injection with other aminoglycosides has not been established.

DOSEAGE FORMS AND STRENGTHS

Piperacillin and tazobactam for injection, USP is supplied as a white to off-white powder.
Each pharmacy bulk package bottle contains piperacillin sodium equivalent to 36 grams of piperacillin and tazobactam sodium equivalent to 4.5 grams of tazobactam.

CONTRAINDICATIONS

Piperacillin and tazobactam for injection is contraindicated in patients with a history of allergic reactions to any of the penicillins, cephalosporins, or beta-lactams inhibitors.

WARNINGS AND PRECAUTIONS

Hypersensitivity Adverse Reactions (including anaphylactic/anaphylactoid reactions) (including shock) have been reported in patients receiving piperacillin and tazobactam for injection. Severe cutaneous adverse reactions are more likely to occur in individuals with a history of severe cutaneous adverse reactions. Patients with a history of severe cutaneous adverse reactions are more likely to experience reactions of sensitivity to multiple allergens. Before initiating therapy with piperacillin and tazobactam for injection, careful inquiry should be made concerning previous hypersensitivity reactions to any of the components of piperacillin and tazobactam for injection should be discontinued and appropriate therapy instituted.

5.2 Severe Cutaneous Adverse Reactions

Piperacillin and tazobactam for injection may cause severe cutaneous adverse reactions, such as Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis. If patients develop a skin rash or other signs and symptoms consistent with piperacillin and tazobactam for injection discontinued if lesions progress.

5.3 Hemophagocytic Lymphohistiocytosis

Cases of hemophagocytic lymphohistiocytosis (HLH) have been reported in patients receiving piperacillin and tazobactam for injection. Signs and symptoms of HLH may include fever, rash, neutropenia, and coagulopathy. Hemophagocytosis, if suspected, discontinue piperacillin and tazobactam immediately and initiate appropriate management.

5.4 Hemolytic Anemia

Bleeding manifestations have occurred in some patients receiving both piperacillin and tazobactam, including piperacillin. These reactions have some been associated with abnormal coagulation tests and clotting time, platelet aggregation and prothrombin time, and are more common in patients with renal impairment. If bleeding occurs, piperacillin and tazobactam for injection should be discontinued and appropriate therapy instituted.

Neutropenia in critically ill patients was associated with piperacillin and tazobactam for injection. Administration appears to be reversible and most frequently associated with prolonged administration. Prompt assessment of hematology should be performed, especially with prolonged therapy, i.e., > 21 days (see Adverse Reactions (5.5)).

5.5 Central Nervous System Adverse Reactions

As with other penicillins, piperacillin and tazobactam may cause neurotoxic effects including seizures. Patients receiving higher doses, especially in the presence of renal impairment may be at greater risk for central nervous system adverse reactions. Closely monitor patients with renal impairment or seizure disorders for signs and symptoms of neurotoxicity or seizures (see Adverse Reactions (5.2)).

5.6 Nephrotoxicity in Critically Ill Patients

The use of piperacillin and tazobactam for injection was found to be an independent risk factor for renal failure and was associated with increased mortality in critically ill patients. Patients receiving higher doses of piperacillin and tazobactam for injection should be monitored for signs and symptoms of renal impairment. Patients receiving higher doses of piperacillin and tazobactam for injection should be monitored for signs and symptoms of renal impairment.

5.7 Electrolyte Effects

Piperacillin and tazobactam for injection contain a total of 23.6 mEq (94 mg Na+) (sodium) per gram of piperacillin in the combination product. This should be considered when treating patients requiring restriction of sodium intake. Patients receiving higher doses should be monitored for signs and symptoms of sodium depletion. Patients with potentially low potassium reserves and who are receiving cytotoxic therapy or diuretics.

5.8 Clostridioides difficile-Associated Diarrhea

Clostridioides difficile-associated diarrhea (CDAD) has been reported with use of many antibiologic agents, including piperacillin and tazobactam for injection, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibiologic agents alters the normal flora of the colon leading to overgrowth of C. difficile.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of C. difficile cause more severe illness and are associated with increased mortality. Appropriate antibiotic therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiologic drug use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibiologic agents.

If CDAD is suspected or confirmed, ongoing antibiologic drug use not directed against C. difficile may need to be discontinued. Appropriate culture and testing should be performed. Supportive treatment, including fluid replacement, protein supplementation, and surgical intervention may be necessary. Patients with severe CDAD should be managed in a hospital setting.

5.9 Development of Drug-Resistant Bacteria

Prescribing piperacillin and tazobactam for injection in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of development of drug-resistant bacteria.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:
Hypersensitivity Adverse Reactions (see Warnings and Precautions (5.1))
Severe Cutaneous Adverse Reactions (see Warnings and Precautions (5.6))
Hemophagocytic Lymphohistiocytosis (see Warnings and Precautions (5.3))
Severe Cutaneous Adverse Reactions (see Warnings and Precautions (5.4))
Central Nervous System Adverse Reactions (see Warnings and Precautions (5.5))
Neutropenia in Critically Ill Patients (see Warnings and Precautions (5.5))
Clostridioides difficile-Associated Diarrhea (see Warnings and Precautions (5.8))

6.1 Clinical Trials Experience

Efficacy trials and mortality were conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Clinical Trials in Adult Patients

During the initial clinical investigations, 2,621 patients worldwide were treated with piperacillin and tazobactam for injection in phase 3 trials. In the key North American monotherapy clinical trials (n=818), 58% of patients were treated with piperacillin and tazobactam for injection in combination with amoxicillin. However, in 32% of the patients treated worldwide, piperacillin and tazobactam for injection was administered as monotherapy. Signs and symptoms of adverse events were mild to moderate in severity and transient in nature. However, in 32% of the patients treated worldwide, piperacillin and tazobactam for injection was administered as monotherapy. Signs and symptoms of adverse events were mild to moderate in severity and transient in nature. However, in 32% of the patients treated worldwide, piperacillin and tazobactam for injection was administered as monotherapy. Signs and symptoms of adverse events were mild to moderate in severity and transient in nature.

Table 6: Adverse Reactions from Piperacillin and Tazobactam for Injection Monotherapy Clinical Trials

System Organ Class
Adverse Reaction
Gastrointestinal disorders
Constipation (7.7%)
Nausea (6.9%)
Vomiting (6.2%)
Diarrhea (5.8%)
Dyspepsia (3.3%)
Flatulence (2.3%)
Abdominal pain (2.3%)
General disorders and administration site conditions
Fever (2.4%)
Injection site reaction (< 1%)
Rigors (< 1%)
Immune system disorders
Allergic reaction (< 1%)
Infections and infestations
Candidiasis (1.6%)
Vaginal candidiasis (1.6%)
Clostridioides difficile-associated diarrhea (1.6%)
Metabolism and nutrition disorders
Hypoglycemia (< 1%)
Musculoskeletal and connective tissue disorders
Myalgia (1.6%)
Arthralgia (< 1%)
Neurological disorders
Headache (7.7%)
Psychiatric disorders
Insomnia (6.2%)
Skin and subcutaneous tissue disorders
Rash (2.4%, including maculopapular, bullous, and urticarial)
Pruritus (2.4%)
Purpura (< 1%)
Vascular disorders
Phlebitis (1.6%)
Thrombophlebitis (< 1%)
Hypotension (< 1%)
Flushing (< 1%)
Respiratory, thoracic and mediastinal disorders
Epistaxis (< 1%)
Nosocomial pneumonia
Two (0.08%) nosocomial respiratory tract infections were conducted. In one study, 222 patients were treated with piperacillin and tazobactam for injection in combination with amoxicillin in combination with ampicillin and 215 patients were treated with ampicillin/clavulanic acid in combination with ampicillin/clavulanic acid. Twenty-five (11%) patients in the piperacillin and tazobactam group and 14 (6.5%) in the ampicillin/clavulanic acid group (> 0.05) also received amoxicillin in combination with ampicillin.

